Pregnant Mar	e Checklist/Records - Mare: _		Foaling Date:	
N	Nare's Udder Filling (2 to 6 wee	ks prior to foaling)	(date)	
N	Mare's Nipples Filling (4-10 days	s prior to foaling)	(date)	
V	ulva/Pelvic Relaxing (up to 3 w	eeks prior to foaling)	(date)	
V	Vaxing/Milk Flow (2-4 days pric	or to foaling)	(date)	
Milk Test Strip: _				(day/results)
Morning Tempera	ture			_ (day/results)
First Stage Labour: (nervous	, swishing tail, discomfort, layi	ng down, sweating, urinatio	on = no more than 2-3	(time/comments hours)
	nr: ntractions increase, sweating, I foot, tie up placenta		sition, appearance of s	(time/comments ack, foot, nose,
Third Stage Labour	(expulsion of placenta wit	hin 3 hours, longer than 3		time/comments)
Weight of Placenta:	Placenta Intact?:	Comments:		

Foal Checklist/Records – Mare:		Birthdate: _	# Days Gestation		
Length of Second Stage Labour:		_ Presentation/Position C	Concerns:		
ime umbilical broke: _	Time Mare got up:	Foal is Alert/Up on	Sternum:Heartrate:		
emperature:	Dipped Umbilical:	Alert:	Attempts to Stand:		
Nursing:	_ Veterinary Exam:	IgG Test:	Pass Meconium:		
After Birth & Within the First 20 Minutes: Moves from laying flat to sternum position, alert & looking about, suckling reflex, possibly moving towards the front of the mare, movement of front legs, reacts to light flash					
Within First 2 Hours: Standing, suckling reflex, stabilization of heart rate & respiratory normal values: respiration less than 40 breaths/min., heart rate less than 80 beats/min, attachment to the mare, nursing					

Within the First 8 Hours: Passing meconium (may be straining), moving about stall, standing up & laying down, nursing, following mare, urinating

Within First 4 Hours: Nursing, moving about the stall, following mare, bright & alert. (If not nursing within the first 4 hours, administer colostrums and call the veterinarian.)

Foal Alert: Watch for behavioural changes. Foal should get stronger each day. Be alert for unusual straining to defacate, nursing too frequently, lethargic/depressed, not laying down easily or not resting enough, diarhea with lethargia/depression, jaundiced appearance to eyes and gums, swelling or lameness of limbs, swelling, discharge or pain of the naval stump.